

Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2024

ミナミマグロ 1 歳魚の加入指標のための
ピストンライン曳縄モニタリング調査 2024 の結果報告

伊藤智幸

Tomoyuki ITOH

国立研究開発法人水産研究・教育機構 水産資源研究所

Fisheries Resources Institute,

Japan Fisheries Research and Education Agency

要約

ミナミマグロ 1 歳魚の加入指数を求める科学曳縄調査を、2024 年 1-2 月に実施した。Esperance から Bremer Bay を含んで Albany までの海域を調査し、ピストンラインは 12 回、調査した。航海を通じて漁獲したミナミマグロは 292 個体であった。40 個体にアーカイバルタグを装着して放流し、ミナミマグロを含む 105 個体の胃内容物を分析し、18 点で CTD による海洋観測を実施した。これまで 1.0 歳魚が主体であったが今回は少なく、今回は初めて 0.3 歳魚が主体となった。

Summary

In January to February 2024, the scientific trolling survey that provides the data for recruitment index of age-1 southern bluefin tuna (SBT) was carried out. The surveyed area extended from Esperance to Albany, including Bremer Bay with 12 times on the piston line. A total of 292 SBT were caught during the survey. 40 SBT individuals were released with archival tags implemented, and the stomach contents of 105 fish including SBT were analyzed. Oceanographic observations were also conducted at 18 locations using CTD. In the past the majority of the fish caught were 1.0 years old, but this time there were fewer, and for the first time, the majority of the fish caught were 0.3 years old.

1. Introduction

Recruitment level is crucial information on stock management of fish, including southern bluefin tuna (*Thunnus maccoyii*, SBT). Several research activities have been attempted for the recruitment monitoring of SBT. Since 1989, Japan has conducted a series of recruitment monitoring survey within a cooperative research framework with Australian scientists. Japan carried out a trolling and pole-and-line catch monitoring survey from 1989 to 1993, and then carried out an acoustic monitoring survey using sonar and echo sounder from 1995 to 2006, for age-1 SBT distributed off the southern coast of Western Australia (Itoh 2006). The recruitment index derived from the acoustic monitoring survey predicted the low recruitment levels of the 1999-2001 year classes of SBT which was confirmed several years later by the model-based assessment so that it was likely to be a reliable index. However, the acoustic survey was ceased after the final survey in 2006 due to the budget restriction. Alternatively, we have carried out a trolling survey since 2006. Australia had carried out a scientific aerial survey in the Great Australian Bight in South Australia since 1993 and provided the recruitment indices, as the aerial survey index, to CCSBT (Eveson et al. 2006). The aerial survey was discontinued after 2017 because of budgetary reason and logistical problems. Since 2016, pilot Gene Tagging (GT) project has been started instead of the aerial survey. GT estimates the absolute amount of stock of age-2 SBT and utilized in the current Management Procedure (MP) (Preece and Bradford 2023).

The trolling survey is a reasonable way to know the recruitment status of age-1 SBT. The recruitment index of age-1 SBT derived from this survey have provided to CCSBT as a fishery independent indicator and utilized in robustness test for MP. This survey finds SBT schools by trolling off the southern coast of Western Australia. Additionally, we set the single straight transect line in the survey area and are investigating intensively each year on this line (Itoh and Kurota 2006). This survey is a long-term survey covering 18 years from 2006 to 2024 except 2015 (Itoh 2021, 2022, 2023, Itoh and Kurota 2006, Itoh and Sakai 2007, 2008, 2009a, 2010, Itoh et al. 2011, 2012a, 2013, Itoh and Tokuda 2014, Itoh and Tsuda 2016, 2020, Tsuda and Itoh 2017a, 2018, 2019). The long time series data can be expected to detect not only the interannual fluctuation of recruitment of age-1 SBT but also a medium-term trend of it. Additionally, the recruitment index from this survey become available immediately in the same year of the survey carried out. Therefore, in order to find recruitment failure, if it occurred, as quick as possible and understand the recruitment trend, it is necessary to continue the trolling survey for the age-1 SBT.

In 2021, due to the global pandemic of COVID-19, the trolling survey was conducted in a limited area in a short period of time. In 2022, the trolling survey was carried out in full range scale including piston-line, though the research items were limited due to still under

the influence of COVID-19. Since 2023, the trolling survey has been able to be carried out in the same specification as before 2020. The trolling index calculated from this survey data is described in another paper (Itoh 2024: CCSBT-ESC/2409/24).

2. Materials and methods

An Australian vessel, *The Southern Conquest* with 17 m in total length, was chartered (Fig. 1). This vessel was consistent to the previous year surveys.

The survey area was off the southern coast of Western Australia between off Esperance (123E) and east of Albany (117E), including off Bremer Bay (Fig. 2). The area covers about 450 km x 60 km. From the coast to offshore the continental shelf of about 70 m in depth extended largely, and then dropped sharply at shelf-edge to deeper than 500 m in depth within 2 km distance. The piston-line laid off Bremer Bay.

The vessels engaged in the survey from 5:30 to 17:00 and returned to a port or a calm bay everyday. GPS position data were recorded every second in a GPS logger device. The vessel operated trolling at speed of 7-8 knots. Eight trolling lines consists of four types of lines at maximum were trolled. Each line has one hook with a plastic lure. Because the trolling index derived from the survey is based on the number of SBT schools, not the total number of SBT individuals caught, we did not try to maximize the number of fish caught. Individual fish caught of any species were measured by its length. Some SBT were killed for biological samples including stomach for contents analysis and mussle tissue for stable isotope analysis. Other SBT were released followed by its fork length (FL) measured. SBT in good condition were tagged with an archival tag (LAT2810L from Lotek Inc.) and released without conventional tags.

Oceanographic observation for vertical profile of temperature and salinity (conductivity) was carried out at 18 points using the Conductivity-Temperature-Depth profiler (JFE Advantech Co. ltd., CTD RINKO-Profiler) to just above sea bottom or 200 m in depth.

3. Results

The trolling survey started on January 30th and finished on February 16th in 2024. After two days survey off Esperance, the vessel moved to Bremer Bay. They surveyed off Bremer Bay area for three days, followed by move west to Albany, and came back to Bremer Bay three days later. They surveyed two days more off Bremer Bay and move east to Esperance. Three more days were surveyed off Esperance. In total, 18 days were surveyed. The piston line were surveyed 12 times.

During the survey, a total of 346 fish were caught, including 292 SBT, 1 skipjack *Katsuwonus pelamis*, 10 yellowtail kingfish *Seriola lalandi*, 11 oriental bonito *Sarda orientalis*, 2 samson fish *Seriola hippos* and others. 91 SBT were killed for stomach contents observation where pilchard *Sardinops sagax* was dominant. Muscle tissue for isotope analysis were sampled from SBT (N=66) and some other species. Total weight of SBT killed was 167.1 kg. 40 SBT were implemented archival tags.

There are four size/age components of SBT in this area at the time of the year: age-0.3 in 30-35 cmFL, age-1.0 having mode at around 48 cmFL, age-1.3 in 57-63 cmFL, and age-2 in 65-80 cmFL (Itoh et al. 2012b). There are two sub-cohorts in age-1 presumably due to different spawning peaks exists in October and January/February. SBT caught in 2024 ranged from 31 to 73 cmFL (Fig. 3). Estimated from fork length, age-0.3 fish was dominant, which is an unusual pattern compared to previous years being age-1.0 dominated. Applying a mixed normal distribution to length frequency estimated that age-0.3 was 55.4%, age-1.0 was 30.0%, age-1.3 was 10.5% and age-2 was 4.2%.

Age-0.3 fish are not rare, but this is the first time they have been the most dominant in SBT. Furthermore, while age-0.3 fish have been caught mostly on the continental shelfedge or in the open ocean up until last year, the area where they are caught (distributed) has expanded to coastal areas. On the other hand, age-1.0 fish which used for the trolling index, caught were few even though a wide area surveyed. Up until now, many age-1.0 SBT have been caught around lumps, but this year, despite persistent and thorough surveys of lumps, no age-1.0 fish were caught.

4. Discussion

Fishery data are basic information to assess the current stock status of SBT in CCSBT. Especially, CPUE of Japanese longline, which covers wide area, season and SBT age range that based on detailed information reported from fishermen, is a long time series index exceeding 50 years used for SBT stock status. The aerial survey was also valuable research. It covered a wide area in a short duration of time in the Great Australian Bight (GAB) by using airplane and data of school biomass is derived as an estimation of a spotter, though ceased in 2017. Gene Tagging is expected to be a fishery-independent index for age-2 SBT and is also used in the Management Procedure. However, there is no single index that can directly reflects the status of SBT stock in whole ranges in both age and geographical distribution. Therefore, we need to collect a variety of information as many as possible to assess the stock status appropriately. Trolling survey provides important information of age-1 SBT abundance.

The design of the trolling survey corresponds to the temporal and spatial distribution

of age-1 SBT, although its survey period and geographic area are limited. A majority of age-1 SBT is thought to be distributed in the coastal area of Western Australia in austral summer (Itoh and Sakai 2009b). Then the age-1 SBT widely migrates to east and west at the end of the summer, while the majority of the SBT move to the GAB. SBT of age-2 and older are distributed widely in the area between off New Zealand and off Cape, and the fish returns to GAB by the following year is not always. Therefore, the recruitment survey targeting age-1 SBT in the southern coast of Western Australia may provide abundance index that represents whole the stock at age-1 SBT.

To derive accurate year trend in an index of recruitment level, we need to exclude other factors that fluctuate by year. Carrying out the survey in a consistent method for all years is a good way for this. We have not changed the gear specification used, and general research method for 18 years. We changed the survey vessel at the 10th and 12th survey, but the type and size of the vessels were consistent all the survey. The survey area has been consistent for 17 years except 2021 where the main survey area has been limited to off Esperance. Such consistent survey design is expected to facilitate to obtain an index that reflects only for annual SBT recruitment change. Additionally, the trolling is robust survey method against environmental factors, including wind, wave, and swell. There were a few days that suspended the research during the cruise due to rough sea condition in 18 years. We did not find a tendency that SBT were less caught in rough weather condition (see Tsuda and Itoh 2017b).

Combining recruitment monitoring data from over 30 years since the early 1990s with information on past fishermen's experience catching SBT, we now know a lot about the distribution of age-1 SBT along the coast of Western Australia. In 2023, few age-1.0 fish were caught around lumps. In 2024, few age-1.0 fish were caught around lumps again, and the catch locations were few. Instead, age-0.3 SBT were widely distributed from offshore to the coast. These are different situations from past distribution knowledge. Furthermore, although there were few age-1.0 fish in the surveys from 2017 to 2024 and the trolling index (TRG) has been low, the CPUE of longline by the corresponding year class remained high (Takahashi and Itoh 2024: CCSBT-ESC/2409/25). This discrepancy between the indices is a concern. If there are changes in the distribution and migration of age-1 SBT that are different from the past, we should understand the situation well and make changes to the trolling survey design if necessary. Also, if there are changes in the distribution and migration of age-1 fish, this may have an impact on GT that use age-2 fish, and MP that use it as input data. Caution should continue to be exercised in comparing the various recruitment indices. This may be something that should be discussed at the ESC, including possible changes in the distribution and migration of age-1 fish, or juvenile fish between age 1 and 4, and their impacts on assessment and

management through GT estimates.

Acknowledgment

We thank Mr. Colin and crew of F/V The Southern Conquest and Mr. Totterdell in MIRG for their help of this survey. We thank Australian Fisheries Management Authority (AFMA) given us scientific research permission in the Australian coastal waters (#1005883).

References

- Eveson, P., Bravington, M., and Farley, J. (2006) The aerial survey index of abundance: updated analysis methods and results. CCSBT-ESC/0609/16.
- Itoh, T. (2006) Acoustic index of age one southern bluefin tuna abundance by the acoustic survey in 2005/2006. CCSBT-ESC/0609/37.
- Itoh, T. (2021) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2020/2021. CCSBT-ESC/2108/29.
- Itoh, T. (2022) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2021/2022. CCSBT-ESC/2208/19.
- Itoh, T. (2023) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2023. CCSBT-ESC/2308/19.
- Itoh, T. (2024) Trolling indices for age-1 southern bluefin tuna: update of the grid type trolling index in 2024. CCSBT-ESC/2409/24.
- Itoh, T. and Kurota, H. (2006) Report on the piston-line trolling survey in 2005/2006. CCSBT-ESC/0609/38.
- Itoh, T. and Sakai, O. (2007) Report on the piston-line trolling survey in 2006/2007. CCSBT-ESC/0709/34.
- Itoh, T. and Sakai, O. (2008) Report on the piston-line trolling survey in 2007/2008. CCSBT-ESC/0809/41.
- Itoh, T. and Sakai, O. (2009a) Report on the piston-line trolling survey in 2008/2009. CCSBT-ESC/0909/32.
- Itoh, T. and Sakai, O. (2009b) Distribution of age 0-1 southern bluefin tuna in Western Australia. CCSBT-ESC/0909/35.
- Itoh, T. and Sakai, O. (2010) Report of the piston-line trolling survey in 2009/2010. CCSBT-ESC/1009/25.
- Itoh, T., Fujioka, K. and Sakai, O. (2011) Report of the piston-line trolling survey in 2010/2011. CCSBT-ESC/1107/29.
- Itoh, T., Sakai, O. and Tokuda, D. (2012a) Report of the piston-line trolling survey in

- 2011/2012. CCSBT-ESC/1208/33.
- Itoh, T., O. Sakai, and D. Tokuda (2012b) Sub-cohort structure of southern bluefin tuna in the recruitment monitoring trolling survey in 2012. CCSBT-ESC/1208/39.
- Itoh, T., Sakai, O. and Tokuda, D. (2013) Report of the piston-line trolling survey in 2012/2013. CCSBT-ESC/1309/27.
- Itoh, T. and Tokuda, D. (2014) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2013/2014. CCSBT-ESC/1409/33.
- Itoh, T. and Tsuda, Y. (2016) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2015/2016. CCSBT-ESC/1609/26.
- Itoh, T. and Tsuda, Y. (2020) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2019/2020. CCSBT-ESC/2008/19.
- Preece, AL. and Bradford, R. (2023) Update on the Genetagging program 2023 and RMA request. CCSBT-ESC/2308/09.
- Takahashi, N. and Itoh, T. (2024) Summary of fisheries and scientific survey indicators of southern bluefin tuna stock in 2024. CCSBT-ESC/2409/25.
- Tsuda, Y. and Itoh, T. (2017a) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2016/2017. CCSBT-ESC/1708/22.
- Tsuda, Y. and Itoh, T. (2017b) Standardization of the grid type trolling index by environmental factors. CCSBT-ESC/1708/24.
- Tsuda, Y. and Itoh, T. (2018) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2017/2018. CCSBT-ESC/1809/26.
- Tsuda, Y. and Itoh, T. (2019) Report of the piston-line trolling monitoring survey for the age-1 southern bluefin tuna recruitment index in 2018/2019. CCSBT-ESC/1909/25.



Figure 1 The Southern Conquest, used for the 2024 trolling survey.
 Photo was taken in 2018.




Figure 2 Trajectory of the vessel with location of southern bluefin tuna caught (circle) in the 2024 trolling survey.




Figure 3 Fork length frequency distribution of southern bluefin tuna caught in the 2024 trolling survey.